

Treeprint When emissions turn personal

ESG Report



04



Executive summary

Trees as a solution for climate change

08 solution



Treeprint: converting emissions

34

into trees



40

43 Emission profiles

44 The Treeprint of our consumers

48 Creating a sustainable lifestyle

Greenhouse gas emissions are personal as they ultimately are generated in order to accommodate our way of life.



Planting trees could be a potential



12

When emissions turn personal

15 In and around the house

20 Eating and Drinking

22 Travel and Tourism

29 Clothing and Shopping

32 Fitness and Entertainment



Consumer profiles: who are you?



53

Emissions: the challenge that lies ahead

Authors

Eugene Klerk, eugene.klerk@credit-suisse.com Betty Jiang, betty.jiang@credit-suisse.com Phineas Glover, phineas.glover@credit-suisse.com

Executive summary

The need for the world to address climate change appears obvious when viewed through the lens of the latest Assessment Report released by the IPCC in early August. In order to meet targets set under the 2015 Paris Agreement, greenhouse gas emissions (GHGs) need to be reduced 50% between 2020 and 2030 and reach 'net zero' by 2050. Much of the debate around climate change centres on what governments or companies say or do. However, we believe this ignores the ultimate driver of emissions: the consumer.

> At the end of the day, it is our daily lifestyle that drives emission levels. Whether it is the car we drive, the flight we take, the products that we buy, or the house that we heat, all emissions globally are ultimately generated in order to accommodate consumer behavior and spending patterns. Therefore, the question is what is our emission footprint and what can be done to bring this in line with climate-change-related limits.

> In this report, we aim to make emissions more personal. We do this by providing emission data for a wide range of activities covering eating and drinking, travel and tourism, entertainment, clothing, and domestic activities (in and around the house). Furthermore, we introduce four different consumer lifestyles to help readers benchmark the emission footprint of their own lives. We conclude that all four of our sample consumers generate emissions in excess of the 3,300kg of CO_o equivalent upper limit that is in line with long-term climate change targets. In fact, for three of the four consumers, emissions need to fall more than 80% in order to reach this limit.

Despite our activity-based emission overview, we believe that some readers may struggle to understand what this emission data actually mean. The concept of a Gigaton (Gt) of emissions is likely to be an alien concept to most of us. This risks making the climate change debate a more theoretical exercise which, if true, hampers the chances of successfully reducing carbon emissions to levels that are in line with long-term climate change targets.



Trees store carbon and could provide a

potential solution. In this report, we introduce the concept of 'Treeprint'. This stands for the number of mature trees that are needed in order to offset the emissions associated with a certain activity. We believe our concept of Treeprint will make it easier for readers to understand the environmental impact of their way of life. Once consumers appreciate how many trees are needed to offset their carbon footprint, they can either decide to reduce certain activities or plant the calculated number of trees in order to reduce their overall emission footprint.

Creating a sustainable consumer. Having reviewed the emission intensity of certain activities and lifestyles and calculating how many trees are needed to offset them, we turn the approach upside down. We ask ourselves what lifestyle has an emission footprint that is within the per capita boundaries implied by long-term climate change targets. Figure 36 on page 49 shows a lifestyle that on our estimates generated c2,600kg of CO₂ per year. The Treeprint of this sustainable consumer is c120 trees, based on our calculations.

We hope that this report will help the reader to understand which areas of his or her lifestyle are most environmentally intense. Furthermore, we hope that this report makes it clear that the biggest impact we can make on a personal level to reduce greenhouse gas emissions is through incorporating changes to our way of life and by planting trees.

David Bleustein Global Head of Securities Research

Trees as a solution for climate change

In recent publications, we highlighted the potential that reforestation may provide in fighting climate change. Specifically, in <u>Credit Suisse Research Institute: The global</u> food system - Identifying sustainable solutions, we first showed that a change in diet would allow for a sharp decline in GHG emissions and free up agricultural land that could be used for reforestation.

In Global ESG Research: The ROE of a Tree, we showed that this reforestation could in theory capture up to 80% of today's anthropogenic emissions and that planting trees could actually be a profitable activity for farmers too.

To outline the role that forests play in storing carbon (natural sequestration), we highlight the following statistics.

- Currently, approximately 30% of CO₂ emissions that are emitted each year get stored or captured by the world's forests, even though respiration and decomposition release some of this carbon back into the atmosphere. Some 47% of emissions remain in the atmosphere; however, calculations by the Intergovernmental Panel on Climate Change (IPCC) suggest that this share is set to grow under more extreme global warming scenarios and without aggressive (re)forestation plans.
- In addition to the carbon that is stored in trees directly, we note that carbon is also stored in soil, primarily through the remains of decomposing plant and animal tissue, and other dissolved organic material. Estimates from the Food and Agriculture Organisation (FAO) indicate that 44% of the carbon stored in forests sits in living biomass, whereas c45% resides in the soil.

- When thinking about using trees to capture carbon, we need to realize that carbon storage differs depending on the type of forest. For example, a 2019 study prepared for the 14th session of the UN Forum on Forests cited a study in Science that found that tropical and boreal (arctic) forests are similar in terms of carbon stock density. Temperate forests, on the other hand, have a carbon stock density that is c40% lower per hectare. We note that much uncertainty remains around these figures, not least because of a lack of data around how much carbon is stored in the soil and how deeply this carbon is stored.
- The fact that natural carbon sequestration is a relevant topic in the fight against climate change is also recognized by institutions such as the IPCC. Its so-called SSP1 scenario requires carbon sequestration through land use (e.g., forests) to more than double during the next 30 years. This, according to the IPCC, is to be achieved by increasing the amount of forest land by c200 million hectares, which could be achieved given that they estimate that the amount of land needed for livestock will decline by more than 300 million hectares.







Figure 1: Natural sequestration is needed to get to net-zero according to IPCC calculations

Source: IPCC





Planting trees could be a potential solution

The potential for carbon storage by trees appears obvious; however, deforestation has consistently reduced the amount of forests and trees during the past few decades. Data from the FAO, for example, suggest that the total area globally that is classified as forest has declined by c178 million hectares since 1990.

Against this background, we note that the topic of reforestation is one that we believe needs to be treated with a greater degree of urgency. The good news is that an increasing number of countries are announcing forestation plans. For example, China recently announced reforestation plans for the next five years. It wants to plant 36,000 square kilometres of new forest (greater than the size of Belgium) per year until 2025 in order to increase its total forest coverage area to

24.1% by 2025 from 23.04% in 2020. China's announcement is part of a range of higher-profile reforestation plans that have been announced by various governments during the past few years. For example, the European Union plans to plant three billion trees by 2030 while the US intends to plant 24 billion trees over the next 20 years.

Figure 3: Tree planting targets for key countries

Country/ Region	Target	Target Year	Annual CO ₂ Emissions (Million tonnes)	Total Trees planted (Million)	Annual CO ₂ captured by mature planted trees (Million tonnes)	Annual CO ₂ captured as % of current emissions
China	36,000 square kilometres of new forest a year	2025	10,175	27,000	594	6%
USA	24 billion trees over 30 years (Trillion Trees Act)	2040	5,285	24,000	528	10%
India	Add 10m ha of new forests and forest cover	2030	2,616	15,000	330	13%
Spain	Nearly 4 million ha increase of forest area by 2032	2032	253	6,000	132	52%
EU	Plant 3 billion trees by 2030	2030	3,749	3,000	66	2%
Brazil	Reforesting 2 million hectares of land by 2030	2030	466	3,000	66	14%
UK	UK forestry cover from 13% to 17% (CCC objectives)	2050	370	1,455	32	9%
Australia	One billion plantation trees by 2030	2030	529	1,000	22	4%
New Zealand	Plant one billion trees between 2018 and 2028	2028	37	1,000	22	60%

Source: Credit Suisse estimates, Total trees planted by country and target year from government announcements

Reforestation on a city level

One of the key challenges in addressing climate change is the need to create engagement with the public. Reforestation programs may act as a partial solution here, especially if these are set up on a more local level with the potential direct involvement of citizens. It appears to us that plenty of opportunities exist for local governments to establish reforestation programs. We have reviewed emission levels for some of the largest urban areas globally.

Figure 4 shows how many trees would have to be planted to offset 25% or 50% of current carbon emissions for some of the largest cities in the world. We have also worked these numbers into the size of forest land that would be needed.

would be captured if an area equivalent to 25% of its city area were planted with trees. Santiago, Chile would need 9% of its city area to capture 25% of its emission levels. For a number of other cities though, we find that reforestation would require a very substantial land area. For example, Singapore would need an area 3.75x its city land area in order to capture 25% of its current carbon emissions. Other cities that require a relatively large area for reforestation include Hong Kong and New York. Clearly, these cities emit a large amount of emissions relative to their size, which is likely due to their highly urbanized, high-rise, structure.

The question that is relevant in this case is whether these plans make a meaningful difference. In Figure 3, we show for a number of countries the percentage of annual emissions that would be captured by these newly planted trees, assuming that they store on average 22kg of CO_o per year. We will address these assumptions later in this report. Our calculations suggest that these reforestation plans, while obviously positive, appear to address less than 15% of current annual CO₂ emissions.

What is apparent to us is that, for a number of these larger cities, a meaningful share of carbon emitted could be captured through reforestation. For example, in the case of Beijing, the city with the highest level of Scope 1 emissions based on 2010 data, we find that 25% of its emissions

Figure 4: Offsetting 25% or 50% of Scope 1 emissions for some of the world's largest cities

							at 50% emi	ssions	at 25% emi	ssions
City	Country	Scope 1 Emissions (Million metric tonnes CO ₂ e)	Emission year	Land area (in square km)	Population (million)	Scope 1 Emissions per Capita	Trees needed (in millions)	Land area required (as % of total land area)	Trees needed (in millions)	Land area required (as % of total land area)
Beijing	China	75.1	2010	16807.8	20.9	3.59	1,707	51%	853	25%
Singapore	Singapore	48.1	2012	728.6	6.0	8.03	1,093	750%	547	375%
Shanghai	China	47.5	2010	6340	27.8	1.71	1,080	85%	540	43%
Hong Kong	China	41.1	2015	1104	7.6	5.41	935	423%	467	212%
New York City	USA	39.0	2015	784	8.2	4.73	886	565%	443	282%
Istanbul	Turkey	37.2	2010	5343	15.4	2.41	845	79%	423	40%
Bangkok	Thailand	27.3	2013	1569	10.7	2.55	620	198%	310	99%
Lagos	Nigeria	25.6	2015	3577	14.9	1.73	583	81%	291	41%
Tokyo	Japan	25.2	2017	2188	37.3	0.67	572	131%	286	65%
Santiago	Chile	23.1	2013	15403	6.8	3.39	525	17%	263	9%
Seoul	South Korea	22.6	2015	605	10.0	2.27	513	424%	257	212%
Mexico City	Mexico	21.3	2016	1485	21.9	0.97	485	163%	242	82%
London	United Kingdom	20.6	2013	1572	9.4	2.19	469	149%	234	75%
Los Angeles	USA	19.6	2013	1215	4.0	4.92	445	183%	223	92%
Houston	USA	19.3	2014	1625	2.3	8.29	438	135%	219	67%
Chicago	USA	16.8	2015	606	2.7	6.27	381	315%	191	157%
Buenos Aires	Argentina	16.3	2015	202	45.2	0.36	371	919%	186	460%
Rio de Janeiro	Brazil	16.2	2012	1224.6	13.5	1.19	368	150%	184	75%
Toronto	Canada	16.2	2013	634	6.3	2.58	367	289%	184	145%
Sao Paulo	Brazil	15.9	2011	1521	22.2	0.71	361	119%	180	59%
Johannesburg	South Africa	15.8	2014	1645	5.9	2.66	358	109%	179	54%
Montreal	Canada	13.6	2009	500	4.2	3.21	310	310%	155	155%
Las Vegas	USA	11.0	2014	136	0.7	16.48	250	919%	125	460%
Bogota	Colombia	10.7	2015	1776	11.2	0.96	243	68%	121	34%
Manchester	United Kingdom	9.6	2013	1277	2.8	3.49	218	85%	109	43%

Source: Citycarbonfootprints.info, CDP, worldpopulationreview.com, Credit Suisse estimates

10008

NAMES AND ADDRESS OF TAXABLE PARTY. or other Designation of the local distance o and the rest of the local division of the lo

NAMES AND ADDRESS OF TAXABLE PARTY.

NAMES OF TAXABLE PARTY.

OTHER DESIGNATION. of the local distance NAMES OF TAXABLE PARTY.





When emissions turn personal

The climate change debate typically focuses on the need for governments to incorporate emission targets and for corporates to cut their emission levels. While this is undoubtedly relevant, it does avoid a focus on the ultimate reason for emissions: the consumer.



Direct residential energy consumption accounts for 22% of total final energy consumption according to the International Energy Agency (IEA). However, one could argue that at the end of the day it is our daily lifestyle that drives energy consumption and by implication emission levels from other sources. Whether it is the car we drive, the products that we buy, or the house that we heat, all emissions globally ultimately are generated by manufacturers or service providers in order to accommodate consumer behavior and spending patterns. The question, therefore, is what is our emission footprint and how can this be offset?

The current distribution of GHG emissions is very uneven across and within countries. For example, Chancel & Piketty ("Carbon and Inequality: from Kyoto to Paris", 2015) showed that the top 10% of GHG emitters globally make up c35% of GHG emissions. Otto et al ("Shift the focus from the super-poor to the super-rich", 2019) showed that the carbon footprint of a typical super-rich household of two is estimated at about 130tCO eq compared to a worldwide average of 3.4tCO eq per capita (Stadler et al, "EXIOBASE 3: developing a time series of detailed environmentally extended multi-regional input-output tables", 2018).

Cutting these per capita emission levels to those that are compliant with per-capita climate change

targets will prove to be a very sizeable challenge. To put this in perspective, on a per capita basis, global emissions need to go down to c2.5-3.3tCO, by 2030, according to analysis from the Global Carbon Project.

Analysis done by Ivanova and Wood in 2020 ("The unequal distribution of household carbon footprints in Europe and its link to sustainability") provides further insight into emission profiles in Europe. They note that only 5% of EU households have emission footprints that are within the targets associated with the Paris Agreement. Strikingly, the top 1% of households by emissions emit c22 times the per capita climate targets (Figure 6).

When reviewing emissions by income category, Ivanova & Wood find that food-related emissions are relatively constant. This does not necessarily mean that people's diets do not change as income changes, but rather that the overall emission profile stays relatively similar.

Areas where emissions appear to change most as income rises include transport, especially air travel. Housing costs are relatively constant except for the top 20% of income levels, where disposable income and savings are likely to have reached sufficiently large sums to increase spending on more emission-intense housing





products and services. In the remainder of this chapter, we will dive deeper into the emission profile of our personal activities in order to provide the reader with a better understanding of how our daily lives and choices impact the environment.



Source: Ivanova, D., & Wood, R. (2020). The unequal distribution of household carbon footprints in Europe and its link to sustainability



Source: Ivanova, D., & Wood, R. (2020). The unequal distribution of household carbon footprints in Europe and its link to sustainability

Our daily lives

We spend our days going through what may often seem like an endless array of activities. Importantly, however, is that all of these activities, from taking a shower in the morning, preparing breakfast, ordering a latte on the way to work or school, having dinner, watching television to using an electric toothbrush, carry an emission footprint.

We believe that the average consumer is likely to underappreciate his or her personal emission footprint. This, in turn, might also help to explain why it appears difficult to get consumers to change their behavior in order to help reduce emissions to achieve climate change targets. With that in mind, we have put together an overview of the emission footprint for a range of ordinary activities. We hope that this will help the reader gain a better understanding of his or her environmental impact.

In and around the house

In our guest to provide readers with insight into their potential personal emission footprint, we first review the carbon intensity of domestic activities or products. Emissions related to energy consumption in buildings have steadily increased. Emissions related to final energy consumption in buildings, for example, have increased by c3% since 1990. However, emissions associated with the upstream power generation needed for this increased by c95%, according to data from the IEA.



8

6

0

Figure 9: Residential building-related energy con





The IEA notes that building sector energy intensity or final energy use per m² of floor space has been declining between 0.5% and 1% per year since 2010. However, this has been more than offset by total floor space growth globally of about 2.5% annually since 2010. Furthermore, what is also apparent is that increased use of more energy efficient sources of building-related energy sources have not managed to replace fossil-fuel-related energy sources.

ssio	ns					
2006	2008	2010	2012	2014	2016	2018
sum	ption	by en	d use			
						

Building-related energy consumption

When reviewing residential energy consumption, we note that space heating and cooling make up the lion's share, with almost 60% on a global basis (Figure 10). Data from EuroStat for the EU

underlines the relevance of heating and cooling in relation to total residential building energy consumption (Figure 11).



Given the high share that space heating and cooling contributes to total residential energy consumption and emission generation, it is vital that this is done in the most efficient way possible. This is even more relevant given that heat generated is lost. Typically, c35% of heat that is lost in buildings is lost through poorly insulated and constructed walls. Some 25% of heat loss occurs through windows and doors while the attic and the basement/floors typically account for 25% and 15% of heat loss, respectively. Overall, there are a number of ways space heating requirements in buildings can be reduced. These methods include:

- 1. Reducing the amount of heat lost through walls and windows through insulation. This includes loft and cavity-wall insulation but may also include insulated wall paper.
- 2. Making the building more airtight and applying heat recovery technologies.
- 3. Installing a more energy efficient heating system.
- 4. Keeping heat below 18°C. Each degree of heating saves around 10% of energy.

Heating systems

Historically speaking, most heating systems use gas or oil as input. However these have a larger carbon footprint than technologies that have been developed more recently. In the chart below, we show the footprint range for a number of these technologies using analysis from the

Figure 11: Carbon footprint range for heating technologies (gram of CO,-eq. /kWh of heat)



Source: The Parliamentary Office of Science and Technology

Parliamentary Office of Science and Technology in the UK and DRAXX. This shows that a substantial reduction in heating-related emission generation can be created by switching to heat pumps and the use of solar-generated heat, particularly when assuming that the energy mix overall continues to switch toward renewables.

Other appliances

In and around our homes, we use a range of devices, each of which creates its own emission footprint. Some of these devices such as washing machines and dishwashers have relatively high carbon footprints; however, one could say that in households these emissions would have to be divided by the number of people making up a family. Personal or private activities around the house that do have a high footprint, in our view, include taking a shower or bath and using the toilet.

In Figure 12, we show some assumptions around the usage of a range of domestic appliances. These range from using the shower for eight minutes each day, using a toilet twice per day, or using an espresso machine four times per day. We calculate the carbon intensity of these activities by multiplying the energy usage or water consumption per minute with the time per year that the appliance is used and then converting this into CO₂ using a conversion factor. This shows that showering and using a bath are relatively emission intense at c280–300kg of CO, per year. Appliances that, when used regularly, also generate sizeable carbon footprints are washing machines, tumble dryers, and dishwashers.

Figure 12: Assumptions around usage of domestic appliances

		(Power) usage	Frequency of use	Duration of use	
Kitchen	Fridge freezer	269kWh per year	Daily	24 hr	
	Dishwasher	1,500 W	1x per day	1 hr	
	Microwave	1,000 W	1x per day	5 min	
	Oven	2,000 W	1x per day	20 min	
	Espresso machine	1,700 W	4x per day	1 min	
	Kettle	2,000 W	4x per day	2min	
	Using the sink	5 litre water/min	3x per day	2 min	
Bathroom	Shower	12 litre/min	1x per day	8 min	
	Toilet	1.6 gallon per flush	2x per day	n.a.	
	Toilet paper	5 sheets per flush	2x per day	n.a.	
	Bath	88 litre per bath	1x per day	n.a.	
Travel	Washing machine	2,000 W	1x per 2 days	1.5 hr	
	Tumble dryer	2,500W	1x per 2 days	1hr	

Source: Credit Suisse estimates



Source: Credit Suisse estimates



Eating and drinking

As we highlighted in our recent Credit Suisse Research Institute report (The global food system), what we eat and drink has a substantial impact in terms of climate change. For the purpose of this report, we highlight the carbon footprint of a range of food and beverage products in Figure 14 and Figure 15.

Meat	Eating a 200g steak						12.00)
	Eating 200g of lamb				8.1	2		
	Eating 200g of chicken	1.50						
Seafood	Eating 200g salmon/tuna	1.58						
	Eating 200g shrimp	2.3	36					
Snacks	Eating 100g of chocolate	0.50						
	Eating 50g of nuts	0.16						
	Eating 50g of crisps	0.12						
Other	Eating 100g of cheese		1.86					
	Eating 2 eggs (100g)	0.42						
	Eating 100g of rice	0.37						
	Eating 100g of greek yogurt	0.28						
	Eating 100g of pasta	0.12						
	Eating 2 slices of bread	0.05						
Fruit/	Eating 100g of berries/grapes	0.14						
vegetables	Eating 100g of potatoes	0.05						
	Eating 100g of vegetables	0.05						
	Eating 1 apple (80g)	0.03						
		0	2	4	6	8	10	

Source: Credit Suisse estimates

Food-related emission data as shown in Figure 14 probably reinforces what most of us have heard before. Eating meat has a far greater emission footprint than a more plant-based or vegetarian diet. However, it is worth highlighting that other food items such as chocolate and cheese have a relatively high emission profile too.

Beverages are far less emission intense than food; however, significant differences between them exist (Figure 15). For example, drinking a pint (just over 500ml) of milk generates 1.28kg of CO_o whereas a pint of beer is almost half that at 0.78kg. Consumers worried about their personal carbon footprint might be pleased to learn that a tea with milk has the lowest carbon footprint even when assuming that the amount of water that is boiled for tea tends to be double the actual amount used for making a cup of tea.





While these individual statistics are interesting, they may not necessarily provide most people with a sense of the carbon intensity of their meals. To provide an indication of this, we have put together a few ordinary dinners and calculated their total footprint.



Source: WWF, BBC Good Food, Credit Suisse estimates



For example, a meal consisting of fish and chips with mushy peas has an average CO₂ emission profile of c1.5kg. Having a steak with fries and cherry tomatoes on the other hand generates close to 10kg of CO_a. We also included a vegetarian dish consisting of pasta bolognese with a plant-based meat alternative. The footprint for this meal is the lowest of all, at just over 0.5kg of CO_a. In other words, a switch from a 'steak frites' to the vegetarian bolognese would result in a 94% drop in emissions.

Travel and tourism

Travel has for years been one of the structural growth themes that we track at Credit Suisse. The expanding middle class across emerging markets and the introduction of low-cost airlines created an environment that provided structural growth in airline travel and tourism as the cost of travel fell. Beyond this, though, we note that travel across all transport modes has seen strong growth during the past few decades. The downside of this, however, is that all travel modes come with sizeable carbon footprints especially when one takes a full life cycle analysis of this into account.

A study published in 2013 by Lenzen et al concluded that global tourism contributed c8% to greenhouse gas emissions (see: <u>The carbon</u> footprint of global tourism | Nature Climate <u>Change</u>). By breaking this down into the various components, the study found that almost 50% of these emissions related to the different transport modes that were used. Other areas that contributed significantly to the overall footprint included consumer goods (12%), food & beverage (10%) and agriculture (8%) (Figure 17). In 2019 and as part of COP25 in Madrid, the World Tourism Organisation (UNWTO) published its findings in relation to transport-related CO emissions of the tourism sector (see: Transportrelated CO, Emissions of the Tourism Sector). It predicted that the number of tourist trips globally was expected to reach more than 37bn by 2030, up 85% from c20bn in 2016. Total transportrelated tourism emissions were forecast by the UNWTO to reach close to 2Gt of CO, which would represent 5.3% of the overall forecast man-made emissions that were forecast by the IEA in the current policies scenario. This compares to 5% in 2016 and 3.7% in 2005. The growth in tourism trips outweighs the benefits from more fuel efficient transport modes.



Source: Nature Climate Change (2018)







Figure 18: Transport-related emissions from tourist arrivals by mode of transport (Mt of CO_2)

Based on the UNWTO calculations as shown in Figure 18, we draw two conclusions. Firstly, and perhaps not surprisingly, air-related emissions make up the majority of international travel. These look set to increase by c50% between 2016 and 2030. Perhaps more important is the rapid growth seen in domestic-related tourism, which we believe is largely a function of the developing middle class across emerging countries. These trips, however, are often undertaken by using cars as transport mode. By 2030 the UNWTO believes that car-related emissions will make up 56% of all domestic travel-related transport emissions and almost 40% of all tourist trips, including international ones. In order to assess how consumers can make a difference, it is therefore important to know how emission intensity differs among transport modes.

The data in Figure 19 show that long-haul flights (especially when sitting in business or first class) are the most emission-intense travel modes. The least intense are, perhaps not surprisingly, trips taken by bicycle or an e-bike.



Tourism not only contributes to emissions via the travel that one needs to undertake to get to a destination but also during the stay at a hotel or resort. With this in mind, we show in Figure 20 the average carbon footprint for hotels based on their star-ranking. We use the Cornell Hotel Sustainability Benchmarking Index for this. The data show that staying one night at a two-star hotel has a median carbon footprint of c15kg of CO_2 . For a five-star hotel, however, the emission intensity increases nearly fivefold to almost 80kg of CO_2 .



el mo	I mode (grams of CO ₂)									
				500						
			121							
	0.45		434							
	245									
	240									
2	234									
212	2									
192										
	200	300	400	500	600					
	_ , ,	200								







Source: Cornell Hotel Sustainability Benchmarking Index

From an environmental perspective, it matters a lot where and how you go on holiday. Data from Cornell's hotel sustainability index show that the average hotel or resort in the Maldives generates more than 160kg of CO_o per night. In Costa Rica and Switzerland, on the other hand, this is less than 10kg of CO_o per night (Figure 21).

In order to make the emission intensity of travel and tourism more understandable, we have put together different types of holidays and calculated their emission profiles. These holidays range from budget holidays in two-star hotels to luxury stays at a resort and a cruise holiday.

By incorporating travel distances to the various destinations and the emission intensity per

kilometre for the various transport modes assumed, we can calculate the combined total emissions for these holidays. Our results show that a true luxury two-week stay at a resort could have an emission footprint that is almost 100x that of a one-week budget holiday taken domestically. Cruise holidays are often seen to be highly emission intense. Our calculations for our cruise holiday suggest that emissions generated while on board are substantially higher than if we were to stay at a hotel. However, these cruiserelated emissions would still be lower than those generated when staying at a high-end resort as our calculations for the 'True luxury' holiday suggest.

Figure 22: Four types of holidays: budget to luxury

Holidays	Nights	Destination	Travel mode	Accommodation
Budget holiday	7	UK domestic	Train	2-star hotel
Mid-budget	7	London - Nice	Air, Economy class	3-star hotel
High-end	14	London - Los Angeles	Air, Business class	4-star hotel
True luxury	14	London - Maldives	Air, First class	5-star resort
Cruise holidays	7	London - Miami	Air, Premium economy	"Cruise liner (3622km round trip)"

Source: Credit Suisse Research

Figure 23: Emission profile for our holidays (kg CO₂)



Travel Accommodation

Source: Credit Suisse estimates

Clothing and shopping

Shopping, particularly fashion, is another highly relevant area of activities that we need to consider when trying to understand our personal emission footprint. The reason for this is that estimates from various sources, such as the UN, McKinsey, and the Ellen MacArthur Foundation, put the contribution of the fashion industry to global greenhouse gas emissions at up to 10%. To put this into context, European Parliamentary Research Service (EPRS) and the UN estimate that the fashion industry's emission intensity is greater than that of all international flights and maritime shipping combined.

Fashion has a sizeable carbon footprint, is often associated with social and labor-related issues. and has a significant environmental impact in terms of water and pesticide usage. For example, the United Nations Environment Programme (UNEP) estimates that the fashion industry is the second-biggest consumer of water and produces 20% of global wastewater.

Figure 24: GHG emissions (upstream) of EU-28 households (2017, indexed to textile/clothing consumption)



Source: Netherlands Organisation for Applied Scientific Research, EU

Figure 24 shows for the EU-28 countries how the greenhouse gas emission profiles for the various activity areas differ. Importantly, this chart also shows what share of these emissions are generated locally and what share is imported because products are produced elsewhere. The chart highlights two important aspects in our view. Firstly, it underlines the relevance of clothing as an emission-intense activity for households. Secondly, it shows that clothing is the activity for which emissions are highly global in nature.

Figure 25: Emission intensity of one pair of Levi's 501 jeans throughout its life cycle is 33.4kg of CO, equivalent



Fashion's footprint: not wearing clothes purchased worsens the issue

A wide range of detailed studies have explored the footprint of the broader clothing or fashion industry. Levi Strauss in 2015 produced a detailed life cycle assessment of its 501 jeans, which we believe provides a clear insight into the drivers behind the intensity of fashion more broadly. The company found that one pair of its 501 jeans is responsible for 33kg of CO emissions throughout its life. In 2018, more than 4.5 billion pairs of jeans were sold globally. Using Levi Strauss's analysis, we calculate that one year of jeans sales alone contribute c150 million tonnes of CO₂ to global greenhouse gas emissions. Other studies that provide insight into the emission footprint of the fashion industry include those produced by WARP in the UK (A Carbon Footprint for UK Clothing and opportunities for savings, 2012) and Roos et al (Environmental Assessment of Swedish fashion consumption, 2015). The conclusions of these studies are in line with the work done by Levi's and show that a significant driver to overall emission intensity relates to the post-production phase (e.g., washing and drying).

However, of equal relevance is the fact that a significant share of clothes are hardly or ever worn, suggesting that all emissions associated with the production, transportation, and retail of these were for nothing. For example, analysis from Weight Watchers in the UK (weightwatchers) showed that 55% of the clothes in an average woman's wardrobe and 47% of clothes in a man's wardrobe are never worn. A survey done by VoucherCloud among women in the US in 2017 found that c20% of the respondents' wardrobes were never worn.

Shopping online is not helping fashion's footprint

One other aspect that is worth mentioning relates to the impact of online versus offline shopping on the clothing footprint. Analysis by MIT's Center for Transportation and Logistics ("Environmental Analysis of US Online Shopping") showed that online shopping has a slightly better emission footprint than when consumers shop in-store (Figure 26). The carbon footprint of a website is smaller than that of a store, whereas parcel carriers are likely to use a more efficient delivery system than a consumer that drives to a store.

However, this conclusion changes when the buying behavior of a 'rushed consumer' is reviewed. Consumers that use 'same-day delivery' options or that order multiple pieces of the same item with the idea to return all but one put significant stress on the delivery system, which significantly increases the emission footprint of the item. In fact MIT's conclusion was that this type of online behavior was less environmentally friendly than if a consumer were to buy in-store.



In Figure 27, we provide carbon footprint data for a range of clothing items. Our analysis suggests that estimates differ even for the same type of item. On average, however, it appears that larger clothing items tend to have a carbon intensity of 10–15kg CO₂ equivalent. Smaller items such as short-sleeved T-shirts emit less than 5kg of CO equivalent during their lifespan.

Solutions to reducing the carbon footprint of clothing appear obvious, as these include investing in higher-guality clothing that lasts longer, wearing items more often, washing them less frequently and at lower temperatures, and not using tumble dryers but air-drying clothes when possible. A life cycle emission footprint would further improve if consumers opted for in-store shopping, especially when using public transport, walking, or cycling as ways to get to their shops.

Figure 27: Carbon footprint for a range of clothing (kg CO,-e)



Source: Levi Strauss, WRAP, Roos et al (2015), Jungmichel (2010), Adidas, Mistra Future Fashion, KTH Royal Institute of Technology

Gym Treadmill (8mph, 30min/d) 44 Entertainment Video gaming (light): High-end PC 41 Streaming (Netflix) on a 50" LED HD TV (2hr/d) 49 Video gaming (extreme): High-end PC Video gaming (extreme): PS4 Pro Video gaming (light): High-end PC OLED TV 60" (3hr/d) 48 Using electronics OLED TV 40" (3hr/d) 42 LED TV 60" (3hr/d) 40 LED TV 40" (3hr/d) 22 Broadband router 36 Desktop PC (8hr use, rest idle) Laptop (8hr use, rest idle) 67 Charging Ipad Air (4th gen.) (1hr/d) 3 electronics Samsung Galaxy (2hr/d) 7 Iphone 13 (2hr/d) 7

Source: Company data, IEA, batteryequivalents.com, Mills et al (2014), Greening the Beast, Credit Suisse Research

Fitness and entertainment

Food, travel, and fashion are activities that receive a lot of attention in relation to emission intensity; however, they are not the only ones that need to be considered, in our view.

Entertainment is one area that might be overlooked by consumers as being emission intense, whether it is watching television, playing videogames, using streaming services, charging a mobile phone or tablet, running on a treadmill or using an exercise bike at the gym. All of this has a carbon footprint. In Figure 28, we summarize what these daily activities contribute to emissions.

Relative to other entertainment activities, we find that videogaming is highly emission intense. An 'extreme' gamer using a high-end PC might be generating more than 450kg of CO, per year. This compares to the 49kg of CO, that is generated by someone who streams a Netflix series for two hours each day on a 50" LED television.

Going to the gym is arguably a positive idea from a health perspective. However, if one also cares about the environment, then it stands to reason that running outside is preferred, given that we estimate a treadmill generates some 44kg of CO₂ per year when used for 30 minutes each day at a speed of 8 miles per hour.







The previous chapter outlined the emission profile for a wide range of activities that consumers are likely to undertake in the course of their daily lives. Based on this, one could draw the conclusion that in order to address climate change we would all simply need to start drinking plant, say, 100 trees to offset the emission tea, not go on holiday anymore, read a book rather than use consumer electronics, wash less frequently, and become a vegetarian. While we see some merit in these activities, individually, we believe it is unrealistic to assume that the majority of consumers globally will adopt such a lifestyle. The question, therefore, is whether something else can be done to reduce the total net impact of a consumer's lifestyle on the environment. We believe that reforestation programs can be an answer or at least part of a solution to this question.

Trees help to explain our emission footprint our **The ROE of a Tree** report, we highlighted

The ability that trees have to store carbon is arguably a good reason for introducing reforestation as a topic in relation to climate change. However, we believe that the concept of trees can also help in making the topic of climate change and the impact of one's personal lifestyle maturity for hundreds of different tree types (see: on it more real and less academic. It likely remains difficult for most consumers to grasp how bad an individual activity is from a climate impact point of view. In other words, we believe that most people do not know how to interpret a certain number of kilograms of CO₀. In order to make it easier for readers to

converting emissions into trees

understand the climate impact of individual activities, we introduce the concept of Treeprint, or the number of mature trees that are needed to offset the emissions generated with a certain activity. Knowing that one has to intensity of a certain activity is, in our view, more likely to resonate with the average consumer than saying that this activity generates 1,000 kilograms of CO_o. Trees, therefore, function both as a conceptual solution as much as a real-life answer to reducing climate change.

We recognize that estimating how many trees are needed to offset a certain level of emissions is not an exact science. A great number of variables play a role including the type of tree, the age of the tree, and where and how the tree is planted. In that average carbon storage for mature trees could be more than 20kg of CO_o per year. Work from the Barcham nursery in the UK is helpful here as it provides very granular data in terms of life expectancy both in rural and urban environments and total carbon storage potential at Barcham). Figure 29 provides implied annual storage numbers for just some of these. Based on Figure 29 alone, we conclude that our assumption that average carbon storage for a tree of 20-25kg per year is achievable.





Based on this data, we have recalculated the emission intensity of the activities highlighted on the previous pages into the number of mature trees that are needed to offset these emissions. In order to indicate the impact of planting trees in different climates, we show the number of trees that are needed for each activity using three different trees that can be planted in a temperate, boreal or, tropical climate. For example, on our analysis, a steak and chips dinner requires 12 mature Birch trees in a temperate climate to offset the related emission footprint. Taking an eight-minute shower each day requires up to ten trees, whereas that luxury holiday would need up to almost 400 trees. If you have two lattes per day, then you need 13 tropical trees to go net zero on this.

Figure 30: The Treeprint of individual activities

Category	Activity	Emis- sions	unit
Eating and drin	king		
Meat	Eating a 200g steak	12.0	kg $\rm CO_2$
	Eating 200g of lamb	8.1	kg $\rm CO_2$
	Eating 200g of chicken	1.5	$\rm kg~CO_{_2}$
Seafood	Eating 200g salmon/tuna	1.6	kg $\rm CO_2$
	Eating 200g shrimp	2.4	kg $\rm CO_2$
Snacks	Eating 100g of chocolate	0.5	kg $\rm CO_2$
	Eating 50g of nuts	0.2	kg CO ₂
	Eating 50g of crisps	0.1	kg CO ₂
Other	Eating 100g of cheese	1.9	kg CO ₂
	Eating 2 eggs (100g)	0.4	kg CO ₂
	Eating 100g of rice	0.4	kg CO ₂
Fruit/vegetable	s Eating 100g of berries/grapes	0.1	kg CO,
	Eating 100g of potatoes	0.1	kg CO
	Eating 100g of vegetables	0.1	kg CO ₂
	Eating 1 apple (80g)	0.0	kg CO
Beverages	Drinking a pint of milk	1.3	kg CO.
J • •	Drinking a pint of beer	0.8	kg CO.
	1 Latte	0.6	ka CO
		0.4	ka CO
	1 Flat white	0.3	ka CO
	1 Espresso (18g coffee)	0.3	kg CO ₂
	Drinking a 150ml glass of wine	0.0	kg CO ₂
	Drinking a cap of regular soda	0.2	kg CO
	Drinking 11 of bottled water	0.2	kg CO ₂
	Drinking in or bottled water	0.2	kg CO ₂
		0.2	kg CO ₂
One person	Pasta bolognese with plant-based	0.1	kg CO ₂
One person dinner	mince		
	Fish and chips with mushy peas	1.5	$\rm kg~CO_{_2}$
	Chicken tikka masala with rice and naan bread	2.0	kg CO ₂
	Ploughman's cheese	2.6	kg $\rm CO_2$
	Lamb stew	5.9	kg CO_2
	Steak and chips with cherry tomatoes	9.6	kg CO ₂
Entertainment a	and related activities		
Charging electronics	Iphone 13 (2hr/d)	7.49	kg CO ₂
	Samsung Galaxy (2hr/d)	7.49	kg CO ₂
	lpad Air (4th gen.) (1hr/d)	3.00	kg $\rm CO_2$
Using	Laptop (8hr use, rest idle)	67.29	kg CO ₂
electronics	Desktop PC (8hr use, rest idle)	123.10	kg CO ₂
	Broadband router	35.95	kg CO,
	LED TV 40" (3hr/d)	22.47	kg CO,
	LED TV 60" (3hr/d)	39.54	kg CO,
			ka CO
	OLED TV 40" (3hr/d)	32.35	NY 60-
	OLED TV 40" (3hr/d) OLED TV 60" (3hr/d)	32.35 48.08	kg CO ₂
Entertainment	OLED TV 40" (3hr/d) OLED TV 60" (3hr/d) Video gamina (light): PS4 Pro	32.35 48.08 40.62	kg CO ₂ kg CO ₂ ka CO
Entertainment	OLED TV 40" (3hr/d) OLED TV 60" (3hr/d) Video gaming (light): PS4 Pro	32.35 48.08 40.62 153.05	kg CO ₂ kg CO ₂ kg CO ₂
Entertainment	OLED TV 40" (3hr/d) OLED TV 60" (3hr/d) Video gaming (light): PS4 Pro Video gaming (light): High-end PC Video gaming (extreme): PS4 Pro	32.35 48.08 40.62 153.05 163.31	kg CO_2 kg CO_2 kg CO_2 kg CO_2
Entertainment	OLED TV 40" (3hr/d) OLED TV 60" (3hr/d) Video gaming (light): PS4 Pro Video gaming (light): High-end PC Video gaming (extreme): PS4 Pro	32.35 48.08 40.62 153.05 163.31 451.79	$kg CO_{2}$ $kg CO_{2}$ $kg CO_{2}$ $kg CO_{2}$ $kg CO_{2}$ $kg CO_{2}$
Entertainment	OLED TV 40" (3hr/d) OLED TV 60" (3hr/d) Video gaming (light): PS4 Pro Video gaming (light): High-end PC Video gaming (extreme): PS4 Pro Video gaming (extreme): High-end PC	32.35 48.08 40.62 153.05 163.31 451.78	$\frac{\text{kg CO}_2}{\text{kg CO}_2}$ $\frac{\text{kg CO}_2}{\text{kg CO}_2}$ $\frac{\text{kg CO}_2}{\text{kg CO}_2}$ $\frac{\text{kg CO}_2}{\text{kg CO}_2}$
Entertainment	OLED TV 40" (3hr/d) OLED TV 60" (3hr/d) Video gaming (light): PS4 Pro Video gaming (light): High-end PC Video gaming (extreme): PS4 Pro Video gaming (extreme): High-end PC Streaming (Netflix) on a 50" LED HD TV (2hr/d)	32.35 48.08 40.62 153.05 163.31 451.78 48.67	$\begin{array}{c} \text{kg CO}_2 \\ \text{kg CO}_2 \end{array}$



		Trees to offset annual emissions by forest type					
Assumptions	Annual CO ₂ emissions (in kgCO ₂)	Temperate - Birch tree (43kg CO ₂ /year)	Boreal - Spruce tree (36kg CO ₂ /year)	Tropical - Eucalyptus tree (32kg CO ₂ /year)			
2 times a weak	1070	11	50	50			
2 times a week	10/2	20	26	10			
2 times a week	024	6	7	40 			
Twice a week	164	4	5	6			
Twice a week	104	3	4	4			
1 ner week	26	1	1	1			
2 times per week	17	1	1	1			
3 times per week	19	1	1	1			
100g per week	97	3	3	4			
Four times a week	87	3	3	3			
Three times a week	58	2	2	2			
2 times per week	15	1	1	1			
2 times per week	5	1	1	1			
4 times per week	10	1	1	1			
4 times per week	7	1	1	1			
1 per dav	466	11	13	15			
1 per dav	285	7	8	9			
2 per dav	402	10	12	13			
2 per day	299	7	9	10			
2 per dav	248	6	7	8			
2 per day	204	5	6	7			
1 glass of 150 ml per day	88	3	3	3			
1 per day	62	2	2	2			
Two litres a day	121	3	4	4			
1 per day	55	2	2	2			
2 per day	51	2	2	2			
Once a week	30	1	1	1			
Once a week	80	2	3	3			
Once a week	106	3	3	4			
Once a week	135	4	4	5			
Once a week	308	8	9	10			
Once a week	499	12	14	16			
2hrs a day, once every two days	7	1	1	1			
2hrs a day, once every two days	7	1	1	1			
1hr a day, once a week	3	1	1	1			
8hr use, 250 work-days	67	2	2	3			
8hr use, 250 work-days	123	3	4	4			
All time use, 365 days a year	36	1	1	2			
3 hours a day	22	1	1	1			
3 hours a day	40	1	2	2			
3 hours a day	32	1	1	2			
3 hours a day	48	2	2	2			
Once a week	41	1	2	2			
Once a week	153	4	5	5			
Once a week	163	4	5	6			
Once a week	452	11	13	15			
2 hours a day	49	2	2	2			
30mins/d, five days a week	44	2	2	2			

In and around t	he house							
Kitchen	Fridge freezer (300w)				122	3	4	4
	Dishwasher (1500W)				248	6	7	8
	Microwave (1000W)				14	1	1	1
	Oven (2000W)				110	3	4	4
	Espresso coffee machine				19	1	1	1
	Kettle (2000W)				55	2	2	2
	Lising the sink				96	3	3	1
Bathroom	Shower				308	8	0	10
Badiloom					00	2	2	10
					7	1	1	1
					000	7	0	0
Weehing	Dain				203	6	0	9
wasning	Tumble driver (2000W)				240	5	6	7
Travel and taxui					201	5	0	1
Transportation	Biovelo	5.0	a CO, por km	5 kilomotros	5	1	1	1
Transportation		5.0	travelled	per day for 200 days	5	1	1	1
	e-bike	7.0	g CO ₂ per km travelled	5 kilometres per day for 100 days	4	1	1	1
	Coach	27.8	g CO ₂ per km travelled	10 kilometres per day for 250 days	69	2	2	3
	Small car (plug-in hybrid electric)	29.4	g CO ₂ per km travelled	10 kilometres per day for 250 days	73	2	3	3
	London Underground	30.8	g CO ₂ per km travelled	10 kilometres per day for 250 days	77	2	3	3
	Light rail and tram	35.1	g CO ₂ per km travelled	10 kilometres for 100 days	35	1	1	2
	National rail	41.2	g CO ₂ per km travelled	1000 kilometres a year	41	1	2	2
	Small electric vehicle (UK electricity)	45.7	g CO ₂ per km travelled	20 kilometres a day for 250 days	183	5	6	6
n e L	Medium electric vehicle (UK electricity)	53.2	g CO ₂ per km travelled	20 kilometres a day for 250 days	213	5	6	7
	Large electric vehicle (UK electricity)	66.9	g CO ₂ per km travelled	20 kilometres a day for 250 days	268	7	8	9
	Medium car (plug-in hybrid electric)	70.8	g CO ₂ per km travelled	20 kilometres a day for 250 days	283	7	8	9
	Bus	104.7	g CO ₂ per km travelled	10 kilometres per day for 250 days	262	7	8	9
	Medium car (hybrid)	109.0	g CO ₂ per km travelled	20 kilometres a day for 250 days	436	11	13	14
	Long-haul flight (economy)	149.8	g CO ₂ per km travelled	2 trips a year (London - New York, 5571 km One-way)	3,338	78	93	105
	Short-haul flight (economy)	155.7	g CO ₂ per km travelled	2 short haul trips a year (London - Paris, 345 km One-way)	215	5	6	7
	Medium car (petrol)	192.3	g CO ₂ per km travelled	20 kilometres a day for 250 days	769	18	22	25
	Black cab (taxi)	211.8	g CO ₂ per km travelled	10 kilometres a day for 250 days	424	10	12	14
	Short-haul flight (business class)	233.6	g CO ₂ per km travelled	2 short haul trips a year (London - Paris, 345 km One-way)	322	8	9	11
	Long-haul flight (economy+)	239.7	g CO ₂ per km travelled	1 trip a year (London - New York, 5571 km One-way)	2,671	63	75	84
	Cruise	245.0	g CO ₂ per km travelled	1 trip of 3622km	887	21	25	28
	Long-haul flight (business class)	434.5	g CO ₂ per km travelled	2 trips a year (London - New York, 5571 km One-way)	9,682	226	269	303
	Long-haul flight (first class)	599.3	g CO ₂ per km travelled	1 trips a year (London - New York, 5571 km One-way)	13,354	311	371	418
Hotel stay by	One night stay at a 2-star hotel	15.1	kg CO,	7 nights stay a year	106	3	3	4
property rating	One night stay at a 3-star hotel	24.2	kg CO ₂	7 nights stay a year	170	4	5	6
	One night stay at a 4-star hotel	41.8	kg CO ₂	7 nights stay a year	293	7	9	10
	One night stay at a 5-star hotel	76.8	$\rm kg~CO_2$	7 nights stay a year	538	13	15	17

Hotel stay by	Costa Rica	6.6	$\rm kg~CO_{_2}$	7 nights stay a year	46	2	2	2
country	Switzerland	8.0	$kg CO_2$	7 nights stay a year	56	2	2	2
	France	8.5	$kg CO_2$	7 nights stay a year	60	2	2	2
	Brazil	8.7	$kg CO_2$	7 nights stay a year	61	2	2	2
	United Kingdom	11.5	${\rm kg}~{\rm CO}_{_2}$	7 nights stay a year	80	2	3	3
	Spain	11.8	$kg CO_2$	7 nights stay a year	83	2	3	3
	United States	19.1	$\rm kg~CO_{_2}$	7 nights stay a year	134	4	4	5
	Portugal	23.8	$kg CO_2$	7 nights stay a year	167	4	5	6
	Italy	26.0	$\rm kg~CO_{_2}$	7 nights stay a year	182	5	6	6
	Australia	34.1	$kg CO_2$	7 nights stay a year	239	6	7	8
	Mexico	38.8	$kg CO_2$	7 nights stay a year	271	7	8	9
	Vietnam	43.1	$kg CO_2$	7 nights stay a year	302	8	9	10
	Turkey	44.6	$kg CO_2$	7 nights stay a year	312	8	9	10
	Egypt	49.3	$\rm kg~CO_{_2}$	7 nights stay a year	345	9	10	11
	South Africa	55.0	$kg CO_2$	7 nights stay a year	385	9	11	13
	Thailand	55.4	$\rm kg~CO_{_2}$	7 nights stay a year	388	10	11	13
	India	55.5	$kg CO_2$	7 nights stay a year	389	10	11	13
	Japan	58.1	$\rm kg~CO_{_2}$	7 nights stay a year	407	10	12	13
	China	59.4	$kg CO_2$	7 nights stay a year	416	10	12	13
	Malaysia	76.1	${\rm kg}~{\rm CO}_{_2}$	7 nights stay a year	533	13	15	17
	United Arab Emirates	81.0	$kg CO_2$	7 nights stay a year	567	14	16	18
	Indonesia	93.3	$kg CO_2$	7 nights stay a year	653	16	19	21
	Maldives	164.5	$\rm kg~CO_{_2}$	7 nights stay a year	1,151	27	32	36
Holiday	Budget holiday		$\rm kg~CO_2$	7 nights UK domestic budget holiday at 2-star hotel, travel by train	130	4	4	5
	Mid-budget		$\rm kg~CO_2$	7 nights mid-budget holiday in Nice at 3-star hotel, air travel by economy class from London	491	12	14	16
	High-end		$\rm kg~CO_2$	14 nights high-end holiday at 4-star hotel, air travel by business class from London	8185	191	228	256
	True luxury		kg CO ₂	14 nights luxury holiday at 5-star hotel, air travel by first class from London	12505	291	348	391
	Cruise holiday		kg CO ₂	7 nights cruise holiday of 3622km round trip including air travel by premium economy from London to Miami	4308	101	120	135

Source: Credit Suisse estimates

Holiday



Consumer profiles: who are you?





We reviewed the emission intensity of individual activities so that our readers can appreciate how damaging these in isolation might be from an environmental perspective. We hope that this is helpful; however, we appreciate that readers might also want to understand how big their overall footprint is likely to be and may want to understand what they can do to offset such an emission footprint. In this chapter, we aim to provide an answer to both these questions.

Firstly, we introduce four hypothetical consumers, each of whom has a different consumption and, therefore, emission profile. Then we calculate how many mature trees would be needed to offset each consumer's emission profile. Our four different types of consumers are as follows:

- The low travel and online consumer: This consumer is assumed to have a diet that has a below average exposure to more emission-intense products such as red meat and dairy. He or she consumes alcohol, although this is less than the consumption of soda and teas. Travel intensity is low and focused on stays in lower star accommodations. We assume that this consumer is more active online, especially via streaming and gaming, and buys more clothing items per year.
- The upwardly mobile consumer: This consumer is assumed to be developing a career. As a result, we believe that this consumer will have an increased travel intensity privately but importantly also business or work related. We assume that this consumer has a somewhat greater focus on health and wellbeing, which translates into going to the gym more often.
- The high carbon consumer: This consumer is believed to travel extensively, likely work related. As far as spending on food and drink is concerned, we believe that this consumer likes to eat meat more than our previous two consumers. We also believe that this consumer drives a larger petrol car and uses domestic appliances more frequently.
- The travel focused consumer: For our final consumer, we assume that his or her lifestyle is focused on travelling, including luxurious holidays overseas. We believe that this consumer has an above-average intake of meat consumption but a below-average consumption of alcohol. Usage of bathroom facilities is assumed to be above average too.

On the following page, we provide an overview of the intensity with which a broad range of products and services are used by our four types of imaginary consumers. The reason for showing this is to enable our readers to assess how similar their lifestyle is to one of our consumers so that an estimate of their emission profile can be made.

Figure 31: Lifestyles for four hypothetical consumers

Category	Metric	Units	Hypothetical life	style profiles by g	eneration	
calogoly		Kg CO _o per	Low travel online	Upwardly mobile	High carbon	Travel focused
		unit	customer	customer	customer	customer
Eating	Consumption of red meat (times per week, 100g each)	6	1	2	3	4
U U	Consumption of poultry (times per week, 100g each)	0.75	3	3	3	2
	Consumption of seafood (times per week, 100g each)	0.79	1	2	1	1
	Consumption of chocolate per week (g)	0.005	100	50	25	0
	Consumption of crisps per week (g)	0.0024	100	75	50	25
	Consumption of cheese per week (g)	0.0186	10	20	25	25
	Number of eggs eaten per week	0.21	2	4	4	3
	Consumption of bread (times per week, 2 slices each)	0.05	6	3	4	5
	Consumption of Greek yogurt (times per week, 100g each)	0.28	1	4	3	2
	Consumption of vegetables per week (times per week, 100g each)	0.05	4	5	6	6
	Consumption of fruit per week (times per week, 100g each)	0.1	5	3	4	5
	Consumption of rice per week (times per week, 100g each)	0.37	3	3	3	2
	Consumption of pasta per week (times per week, 100g each)	0.12	3	2	2	4
Drinking	Consumption of potatoes per week (times per week, 100g each)	0.05	5	2	2	4
Drinking	Number of beer units per week	0.76	0	1	7	2
	Number of wine units per week	0.24	7	7	2	1
	Number of coffees per day	0.10	1	1	5	3
	Litres of bottled water per day	0.40	0.5	20	1.5	1.0
	Number of teas per day	0.07	4	1	0	5
	Number of units of milk per week	1.28	4	2	2	2
Travel	Number of times taking long-haul return flights per year (husiness	0.434	0	2	4	2
	class) (3500km)	0.101		£		-
	Number of times taking long-haul return flights per year (premier economy)	0.24	0	1	1	0
	Number of times taking short-haul return flights per year (business)	0.234	0	0	2	0
	Number of times taking short-haul return flights per year (economy)	0.156	2	10	15	5
	Average km in a car per year (large petrol)	0.283	0	0	10000	15000
	Average km in a car per year (medium, petrol)	0.192	0	5000	0	0
	Average km in a car per year (mediam) periol)	0.154	1000	0000	0	
	Average km in a car per year (large, electric)	0.067				
	Average km in a car per year (medium, electric)	0.053				
	Average km in a car per year (small, electric)	0.046				
	Number of days using intra city public transport per week		5	5	3	1
	Average distance travelled by public transport each time (km)	0.033	5	20	14	10
	Number of times using long distance rail/bus travel per month		2	1	0	0
	Average distance travelled by long distance rail each time (single, km)	0.0345	200	100		
	Average km cycling per week	0.005	50	50	0	0
Tourism	Average nights spent in a 2-star hotel per vear	15.1	5	00	0	0
louisin	Average nights spent in a 3-star hotel per year	24.2	5			
	Average nights spent in a 4-star hotel per year	41.8		10		5
	Average nights spent in a 5-star hotel per vear	76.8		10	20	10
	Average nights spent in high-end luxury hotel/cruise per vear	145		-		10
Fitness	Number of times used treadmill per week (30min each)	0.12	3	5	3	2
Entertainment	Average time per day spent watching regular broadcast TV (in	0.02	1.0	1.0	1.0	2.5
	Average time per day spent watching streaming TV (e.g., Netflix)	0.066	1.0	2.0	2.0	2.0
	Average time per day spent video gaming (hours)	0.10	2.5	1.0	0.0	0.0
	Use of a broadband router	36	1	1	1	1
	Average time per day using a desktop (hours)	100	3.0	8.0	6.0	1.0
	Average time per day using a laptop (hours)	50	3.0	1.0	1.0	1.0
	Average time per day charging a smartphone (hours)	25	2.0	2.0	2.0	1.0
In and around the	Number of times using washing machine per week (1.5hr wash	1.2	1	2	3	2
	each)	0.0	7	7	6	5
house	Number of times taking a both per week (omin each)	0.0	<i>i</i>	0	1	0
nouse	Number of times flushing a toilet per dev (1.6 college of water cook)	0.0	0	0	3	2
	Number of times hushing a tollet per day (1.0 gallons of Water each)	100	<u> </u>	<u>∠</u> 1	1	4
	Number of times a dishwasher is used per week (1500W/ 1br)	0.6	1	3	5	3
	Number of times a oven is used per week (1000W, 111)	0.3	3	4	5	5
Clothing/fashion	Number of jeans/trousers bought per vear	20	3	3	2	2
ciounig/ lashioli	Number of shirts hought per year	11	4	4	4	3
	Number of jackets bought per year	13.5	2	1	2	0
	Number of suits/dresses bought per vear	15	1	3	2	0
	Number of t-shirts/underwear small items bought per year	3	10	8	6	5

Source: Credit Suisse estimates

Emission profiles

Based on the assumptions as highlighted in Figure 31 and using carbon intensity estimates for each of them, we can calculate the potential annual emission intensity for our four different types of consumers.

Our assumptions suggest that our 'high carbon' consumer generates c25,000kg of CO₂ per year, or roughly 8x the level generated by our 'low travel and online' consumer (Figure 32). The emission profiles of the other two consumers are fairly similar at 15–17 thousand kilograms of CO, per year.

To put these numbers into context, we refer back to page 13, where we highlighted that per capita emissions needed to be 2.5–3.3Gt of CO₂ per year in order to be in line with Paris Agreement related targets. Taking the midpoint of 2.9Gt of CO₂, we calculate that average emissions for our



Figure 32 shows the main factors in terms of emissions for each of our four consumers. The answer to the guestion of how to reduce emissions to sustainable levels appears obvious. Travel intensity needs to reduce or less emissionintense forms of travel or tourism need to be chosen. From an environmental perspective, it makes more sense to take a train and go camping domestically than to fly business class and stay in a five-star hotel far away.

'upwardly mobile', 'high carbon', and 'travel focused' consumers need to fall between 81% and 88%. Even for our 'low travel and online' consumer we note that emission levels are c20% too high.



Figure 33: Share of annual emission intensity of our four consumers by category (kg CO_2 equivalent)

The Treeprint of our four consumers

In addition to cutting or changing behavior in order to reduce one's carbon footprint, consumers can also opt for carbon offsetting schemes such as planting trees. Our data for carbon storage by different types of trees allow us to calculate how many mature trees are needed to offset a certain level of emissions. Based on the emission profiles for our four consumers, we have calculated how many trees they would need in order to arrive at a net-zero state (Figure 34). We also provide the number of trees needed to offset the activity with the greatest carbon footprint for each of our four consumers (Figure 35). Our calculations indicate that consumers who like their meat, go on overseas holidays or travel and take a plane to get there, and that frequently use domestic appliances, including consumer electronics, require 700–1,140 mature trees in order to offset their private carbon footprint. Their Treeprint range is 700-1,140.





Source: Credit Suisse estimates

Figure 35: Offsetting emissions for the activity with the highest contribution to overall emissions



Source: Credit Suisse estimates

Some readers might decide that having a Treeprint of, say, 1,000 trees is great as this suggests that, by planting that many trees, they would apparently offset their personal carbon footprint and be able to live without altering their lifestyle. We highlight the following reasons why such a conclusion could be wrong.

- Firstly, we note that, for a number of activities highlighted previously, we only consider direct emissions rather than the full life cycle impact of that activity. Our Treeprint estimate is, therefore, likely to underappreciate the full impact of an activity or lifestyle.
- Secondly, we note that our calculations are based on the carbon storage potential of mature trees. Planting trees now is obviously a good decision but, in most cases, it will take quite a few years before these trees reach their maximum annual carbon storage potential. During these initial years, a consumer planting trees would therefore not lead a "net-zero lifestyle".

Ŵ

 Thirdly, our Treeprint calculation provides the number of mature trees that are alive and store carbon. Not all trees that are planted today, however, will survive long enough to reach maximum carbon storage potential; therefore, consumers would need to plant more trees than our Treeprint estimate.

Finally, there is a practical limitation to consider. Currently, there are some 5bn people globally between the ages of 20 and 79. Most of these, however, live in developing economies, suggesting that their Treeprint is unlikely to reach anywhere near the levels suggested for our hypothetical consumers. However, if we were to assume that only 20% of all adults have a meaningful Treeprint of, say, 500 trees, we would already be looking at a tree planting requirement of 0.5 trillion trees. That alone represents 16% of the current total number of trees globally of c3trn.

Creating a sustainable lifestyle

Up until now, we have reviewed consumer profiles of different types of consumers, estimated their emission footprint, and concluded that these were not sustainable in that the average emissions generated per person were higher than the limit that is deemed in line with long-term climate change targets. We now turn our quest upside down and ask ourselves what sort of lifestyle is sustainable or has an emission profile of 2,500-3,300kg of CO₂ equivalent per year.

We obviously know that there is no one lifestyle that is sustainable and that personal choice will ultimately determine how one becomes sustainable if that is what one wants to achieve. Below we show one example of a lifestyle that has an emission profile that meets the CO_o boundary condition of less than 3,330 Gt of CO per year. We make the following comments about our assumptions for the three largest areas.

- Eating and Drinking: The key emission drivers here are: i) red meat, ii) beer, iii) milk, and iv) coffee consumption. Our sustainable consumer has swapped these for poultry, sea food, tea, and some wine.
- Travel: As shown before, travel is emission intense. Therefore it stands to reason that living sustainably means limited (long haul) flying and reduced use of large petrol cars. Our sustainable consumer takes one holiday per year consisting of a short-haul economy class flight and a four-star hotel. He or she has opted for an electric rather than petrol car and drives 5,000km per year. The remainder of the time rail, inner-city public transport, and a bicycle are used.

In and around the house: This is a difficult one given that most of the related activities (e.g., washing, cleaning) are those that are most difficult to change. Our sustainable consumer takes six showers per week but for four minutes each rather than the usual eight. The washing machine is used two times per week, however, on a shorter program of 1hr rather than the usual 1.5hr. Finally, the dishwasher is used only once per week as our consumer has opted for hand washing the dishes during the other days.

Overall, we estimate that the lifestyle of our sustainable consumer generates c2.6Gt of CO_o per year, putting it well within the maximum boundary for a sustainable lifestyle. The Treeprint of this consumer is c120 trees, based on our calculations. For some readers, adopting such a lifestyle in order to live sustainably may be difficult to accept; however, we note that in order to make our world more sustainable the a "no-change" scenario is simply unrealistic. We hope that this report will help the reader to identify which areas of his or her lifestyle are most environmentally intense and how many trees need to be planted in order to minimize his or her carbon footprint.









Figure 38: Hypothetical lifestyle for a consumer with a sustainable carbon footprint

Category	Metric	Frequency of activity of the sustainable consumer
Eating	Consumption of red meat (times per week, 100g each)	0
	Consumption of poultry (times per week, 100g each)	2
	Consumption of seafood (times per week, 100g each)	2
	Consumption of chocolate per week (g)	50
	Consumption of crisps per week (g)	50
	Consumption of cheese per week (g)	25
	Number of eggs eaten per week	2
	Consumption of bread (times per week, 2 slices each)	3
	Consumption of greek yogurt (times per week, 100g each)	3
	Consumption of vegetables per week (times per week, 100g each)	7
	Consumption of truit per week (times per week, 100g each)	5
	Consumption of rice per week (times per week, 100g each)	3
	Consumption of pasta per week (times per week, 100g each)	2
	Consumption of potatoes per week (times per week, 100g each)	2
Drinking	Number of beer units per week	0
	Number of wine units per week	4
	Number of (diet) soda cans per week	4
	Number of coffees per day	0
	Litres of bottled water per day	2
	Number of teas per day	4
	Number of units of milk per week	0
Travel	Number of times taking long-haul return flights per year (business class) (3500km)	0
	Number of times taking long-haul return flights per year (premier economy)	0
	Number of times taking short-haul return flights per year (business) (500km)	0
	Number of times taking short-haul return flights per year (economy)	1
	Average km in a car per year (large, petrol)	0
	Average km in a car per year (medium, petrol)	0
	Average km in a car per year (small, petrol)	
	Average km in a car per year (large, electric)	
	Average km in a car per year (medium, electric)	5000
	Average km in a car per year (small, electric)	
	Number of days using intra city public transport per week	4
	Average distance travelled by public transport each time (km)	5
	Number of times using long distance rail/bus travel per month	2
	Average distance travelled by long distance rail each time (single, km)	200
	Average km cycling per week	30
Tourism	Average nights spent in a 2-star hotel per year	
	Average nights spent in a 3-star hotel per year	
	Average nights spent in a 4-star hotel per year	10
	Average nights spent in a 5-star hotel per year	0
	Average nights spent in high-end luxury hotel/cruise per year	0
Fitness	Number of times used treadmill per week (30min each)	3
Entortoinment	Average time per day spent watching regular broadcast TV (in hours)	1
Entertainment	Average time per day spent watching streaming TV (e.g., Netflix) (in hours)	1
	Average time per day spent video gaming (hours)	0
	Use of a broadband router	1
	Average time per day using a desktop (hours)	8
	Average time per day using a laptop (hours)	1
	Average time per day charing a smartphone (hours)	1
In and around the house	Number of times using washing machine per week (1hr wash each)	2
	Number of times using the shower per week (8min each)	3
	Number of times taking a bath per week (88 litres each)	1
	Number of times flushing a toilet per day (1.6 gallons of water each)	2
	Use of a fridge freezer	1
	Number of times a dishwasher is used per week (1500W, 1hr)	1
	Number of times an oven is used per week (2000W, 20min)	3
Clothing/fashion	Number of jeans/trousers bought per year	2
	Number of shirts bought per year	3
	Number of jackets bought per year	0
	Number of suits/dresses bought per year	0
	Number of t-shirts/underwear small items bought per year	5

Source: Credit Suisse estimates

In this report, we addressed the emission profile of more personal activities as this is, in our view, a significant factor in driving overall emissions. With that in mind, it might be helpful to summarize the challenge that the world faces in terms of emissions.

Annual greenhouse gas emissions which include carbon dioxide (CO_2) as well as a number of other gases such as methane (CH_4) and nitrous oxide (N_2O) are currently running at an annual rate of c50Gt of CO_2 equivalent. Carbon dioxide emissions alone have increased sevenfold since the 1950s and almost 10x during the past 100 years (Figure 39).

Ŵ

Emissions: the challenge that lies ahead



Figure 40: Atmospheric concentration of carbon dioxide



The implications of the rise in emissions are becoming ever more visible. Hot extremes (including heatwaves) have become more frequent since the 1950s, whereas cold extremes have become less frequent. Furthermore, the frequency of events such as flooding and hurricanes has also increased, resulting in significant economic and environmental damages and frequent loss of life.

In its latest report, the IPCC also notes that sea levels are likely to rise further as a result of global warming. These, they believe, could rise by 19-22 metres during the next 2,000 years if average temperatures rise by 5°C.

I his s
emiss
could
increa
centu
mear
extre
once
often
by 21
frequ
increa
agric
occui
and e
devel



Source: IPCC, AR6 Page 41



To underline how significant climate change may become, the IPCC has developed five scenarios. shows that, without substantial changes to ssion generation going forward, the world indeed be heading for average temperature ases of more than 5°C by the end of this ury. The IPCC also outlined what this could n for extreme weather events. For example, eme temperature events that currently occur every ten years are likely to occur 9x more if average temperatures increase by c5°C 100 from pre-industrial levels. The uency of extreme rainfall events would ease c3x in this scenario while extreme cultural and ecological drought is likely to ur 4x more often. The environmental, social, economic implications of these elopments cannot by overestimated, in our

view, and suggest that strong action is needed.

The IPCC has clearly outlined what needs to happen to emissions if the world is to minimize average temperature increases. For example, to limit average temperature increases to $c1.5^{\circ}$ C by the end of 2100, total CO₂ emissions need to fall c50% between 2020 and 2030; they then need to reach net zero by 2050 before declining further to reach a net-negative 10 Gt of CO₂ per year by 2100.



Source: IPCC



CREDIT SUISSE LEGAL ENTITY credit-suisse.com

While this post has been prepared by the Securities Research business of Credit Suisse AG, its subsidiary or affiliate ("CS") and may contain references to Securities Research reports and/or Securities Research analysts, it is for information only and does not constitute research by CS. Furthermore, this post is not to be regarded as a sales prospectus or an offer or solicitation of an offer to enter in any investment activity. This post does not take into account your specific investment objectives and needs nor your financial situation. No representation or warranty, either expressed or implied is provided in relation to the accuracy, completeness or reliability of the information contained herein, nor is it intended to be a complete statement or summary of the developments referred to in this post and any liability therefore (including in respect of direct, indirect or consequential loss or damage) is expressly disclaimed. The information and any opinions expressed in this post are subject to change without notice and may differ or be contrary to opinions expressed by other business areas or groups of CS as a result of using different assumptions and criteria. CS is not providing any financial, economic, legal, accounting, or tax advice or recommendations in this post. In addition, the receipt of this post alone by you is not to be taken to constitute you as a client of any CS entity. CS is under no obligation to update or keep current the information contained in this material. Please consult with your client advisor before taking any investment decision. This material is issued and distributed in the United States by CSSU, a member of NYSE, FINRA, SIPC and the NFA, which accepts responsibility for its content. Clients should execute transactions through a Credit Suisse entity in their home jurisdiction unless governing law permits otherwise.

© 2021, CREDIT SUISSE LEGAL ENTITY.

